

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
2 September 2004 (02.09.2004)

PCT

(10) International Publication Number
WO 2004/074289 A1

(51) International Patent Classification⁷: **C07D 487/04**,
A61K 31/495, A61P 1/04

(21) International Application Number:
PCT/EP2004/050135

(22) International Filing Date: 16 February 2004 (16.02.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
03003652.9 18 February 2003 (18.02.2003) EP

GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (for all designated States except US): **ALTANA PHARMA AG** [DE/DE]; Byk-Gulden-Str. 2, 78467 Konstanz (DE).

(72) Inventors: **CHIESA, M.**, Vittoria; Alpenstr. 5, 78464 Konstanz (DE). **PALMER, Andreas**; Pirminweg 1a, 78467 Konstanz (DE). **BREHM, Christof**; Richentalstr. 8, 78462 Konstanz (DE). **GRUNDLER, Gerhard**; Meersburger Str. 4, 78464 Konstanz (DE). **SENN-BILFINGER, Joerg**; Saentisstrasse 7, 78464 Konstanz (DE). **SIMON, Wolfgang-Alexander**; Schubertstrasse 17, 78464 Konstanz (DE). **POSTIUS, Stefan**; Austrasse 4b, 78467 Konstanz (DE). **KROMER, Wolfgang**; Hinterhauserstr. 5, 78464 Konstanz (DE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **ZIMMERMANN, Peter, Jan** [DE/DE]; Zum Lerchental 43/1, 78315 Radolfzell (DE). **BUHR, Wilm** [DE/DE]; Zum Kirchenwald 7, 78465 Konstanz (DE).

(74) Agent: **RUPP, Herbert**; c/o ALTANA Pharma AG, Byk-Gulden-Str. 2, 78467 Konstanz (DE).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,

Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, ARIPO patent (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)
- of inventorship (Rule 4.17(iv)) for US only

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: 6-SUBSTITUTED IMIDAZOPYRAZINES

(57) Abstract: The Invention relates to 6-substituted imidazopyrazines of formula I, in which the substituents and symbols have the meanings indicated in the description. The compounds have gastric secretion inhibiting and excellent gastric and intestinal protective action properties.

WO 2004/074289 A1

1 JC20 Rec'd PCT/PTO 11 AUG 2005

6-Substituted imidazopyrazinesTechnical field


The invention relates to novel compounds, which are used in the pharmaceutical industry as active compounds for the production of medicaments.

Prior art

In European patent applications 204285 (which corresponds to US patents 4,725,601 and 4,782,055), certain imidazoheterocyclic compounds are disclosed, which are said to be useful in the treatment of ulcers. In European patent application 299470 (which corresponds to US patent 5,112,834), certain imidazopyridines and -pyrazines are disclosed, which are said to have an excellent protective effect on the stomach and intestine of warm-blooded animals. International patent application WO 99/28322 (which corresponds to US patent 6,518,270) describes heterocyclic compounds, among others imidazopyrazines with a certain substitution pattern, which are said to inhibit gastric acid secretion. In J. Med. Chem. 1987, 30, 2031 - 2046, J. Kaminski et al. describe the gastric antisecretory, cytoprotective and metabolic properties of certain substituted imidazo[1,2-a]pyrazines, which are unsubstituted in the 6-position. In 'The Practice of Medicinal Chemistry', pages 203 - 237, C. Wermuth gives a review on 'Molecular Variations based on Isosteric Replacements'. In the abstract of Japanese Patent publication No. 07242666, a variety of heterocyclic compounds is disclosed, amongst which imidazo[1,2-a]pyridines are exemplarily named, which are said to be useful for preventing and treating allergy, inflammation, autoimmune diseases, shock, ache, etc., as bradykinine-antagonizing agents. In International patent application WO 02/060492 certain (condensed) pyridine and pyrazine derivatives are disclosed, which are said to be useful as kinase inhibitors.

Summary of the invention

The invention relates to compounds of the formula 1



in which

R1 is hydrogen, 1-4C-alkyl, 3-7C-cycloalkyl, 3-7C-cycloalkyl-1-4C-alkyl, 1-4C-alkoxy, 1-4C-alkoxy-1-4C-alkyl, 1-4C-alkoxycarbonyl, 2-4C-alkenyl, 2-4C-alkynyl, fluoro-1-4C-alkyl or hydroxy-1-4C-alkyl,

R2 is hydrogen, 1-4C-alkyl, aryl, 3-7C-cycloalkyl, 3-7C-cycloalkyl-1-4C-alkyl, 1-4C-alkoxycarbonyl, hydroxy-1-4C-alkyl, fluoro-1-4C-alkyl, halogen, 2-4C-alkenyl, 2-4C-alkynyl, amino, mono- or di-1-4C-alkylamino-1-4C-alkyl or cyanomethyl,

R3 is halogen, fluoro-1-4C-alkyl, 2-4C-alkenyl, 2-4C-alkynyl, carboxyl, cyano, 1-4C-alkoxycarbonyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy-1-4C-alkyl, 1-4C-alkoxy-1-4C-alkoxy-1-4C-alkyl, fluoro-1-4C-alkoxy-1-4C-alkyl or the group -CO-NR31R32,

where

R31 is hydrogen, hydroxyl, 1-7C-alkyl, hydroxy-1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl and

R32 is hydrogen, 1-7C-alkyl, hydroxy-1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl,

or where

R31 and R32 together, including the nitrogen atom to which both are bonded, are a pyrrolidino, piperidino, piperazino, N-1-4C-alkylpiperazino or morpholino group,

X is O (oxygen) or NH and

Ar is a mono- or bicyclic aromatic residue, substituted by R4, R5, R6 and R7, which is selected from the group consisting of phenyl, naphthyl, pyrrolyl, pyrazolyl, imidazolyl, 1,2,3-triazolyl, indolyl, benzimidazolyl, furyl, benzofuryl, thienyl, benzothienyl, thiazolyl, isoxazolyl, pyridinyl, pyrimidinyl, chinolinyl and isochinolinyl,

wherein

R4 is hydrogen, 1-4C-alkyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy, 2-4C-alkenyloxy, 1-4C-alkylcarbonyl, carboxy, 1-4C-alkoxycarbonyl, carboxy-1-4C-alkyl, 1-4C-alkoxycarbonyl-1-4C-alkyl, halogen, hydroxy, aryl, aryl-1-4C-alkyl, aryl-oxy, aryl-1-4C-alkoxy, trifluoromethyl, nitro, amino, mono- or di-1-4C-alkylamino, 1-4C-alkylcarbonylamino, 1-4C-alkoxycarbonylamino, 1-4C-alkoxy-1-4C-alkoxycarbonylamino or sulfonyl,

in which

aryl is phenyl or substituted phenyl with one, two or three identical or different substituents from the group of 1-4C-alkyl, 1-4C-alkoxy, carboxy, 1-4C-alkoxycarbonyl, halogen, trifluoromethyl, nitro, trifluoromethoxy, hydroxy and cyano,

R5 is hydrogen, 1-4C-alkyl, 1-4C-alkoxy, 1-4C-alkoxycarbonyl, halogen, trifluoromethyl or hydroxy,

R6 is hydrogen, 1-4C-alkyl, 1-4C-alkoxy, hydroxy or halogen and

R7 is hydrogen, 1-4C-alkyl or halogen,

and the salts of these compounds.

1-4C-Alkyl represents straight-chain or branched alkyl groups having 1 to 4 carbon atoms. Examples which may be mentioned are the butyl, isobutyl, sec-butyl, tert-butyl, propyl, isopropyl, ethyl and the methyl group.

3-7C-Cycloalkyl represents cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl, of which cyclopropyl, cyclobutyl and cyclopentyl are preferred.

3-7C-Cycloalkyl-1-4C-alkyl represents one of the aforementioned 1-4C-alkyl groups, which is substituted by one of the aforementioned 3-7C-cycloalkyl groups. Examples which may be mentioned are the cyclopropylmethyl, the cyclohexylmethyl and the cyclohexylethyl group.

1-4C-Alkoxy represents a group, which in addition to the oxygen atom contains one of the aforementioned 1-4C-alkyl groups. Examples which may be mentioned are the butoxy, isobutoxy, sec-butoxy, tert-butoxy, propoxy, isopropoxy and preferably the ethoxy and methoxy group.

1-4C-Alkoxy-1-4C-alkyl represents one of the aforementioned 1-4C-alkyl groups, which is substituted by one of the aforementioned 1-4C-alkoxy groups. Examples which may be mentioned are the methoxymethyl, the methoxyethyl group and the butoxyethyl group.

1-4C-Alkoxy-carbonyl (1-4C-alkoxy-C(O)-) represents a carbonyl group, to which one of the aforementioned 1-4C-alkoxy groups is bonded. Examples which may be mentioned are the methoxycarbonyl ($\text{CH}_3\text{O}-\text{C}(\text{O})-$) and the ethoxycarbonyl group ($\text{CH}_3\text{CH}_2\text{O}-\text{C}(\text{O})-$).

2-4C-Alkenyl represents straight-chain or branched alkenyl groups having 2 to 4 carbon atoms. Examples which may be mentioned are the 2-butenyl, 3-butenyl, 1-propenyl and the 2-propenyl group (allyl group).

2-4C-Alkynyl represents straight-chain or branched alkynyl groups having 2 to 4 carbon atoms. Examples which may be mentioned are the 2-butyne, 3-butyne, and preferably the 2-propyne, group (propargyl group).

Fluoro-1-4C-alkyl represents one of the aforementioned 1-4C-alkyl groups, which is substituted by one or more fluorine atoms. An example which may be mentioned is the trifluoromethyl group.

Hydroxy-1-4C-alkyl represents one of the aforementioned 1-4C-alkyl groups, which is substituted by a hydroxy group. Examples which may be mentioned are the hydroxymethyl, the 2-hydroxyethyl and the 3-hydroxypropyl group.

Halogen within the meaning of the invention is bromo, chloro and fluoro.

Mono- or di-1-4C-alkylamino represents an amino group, which is substituted by one or by two - identical or different - groups from the aforementioned 1-4C-alkyl groups. Examples which may be mentioned are the dimethylamino, the diethylamino and the diisopropylamino group.

Mono- or di-1-4C-alkylamino-1-4C-alkyl represents a 1-4C-alkyl group, which is substituted by a mono- or di-1-4C-alkylamino group. Examples which may be mentioned are the dimethylaminomethyl, the diethylaminomethyl, the methylaminomethyl and the diisopropylaminomethyl group.

1-4C-Alkoxy-1-4C-alkoxy represents one of the aforementioned 1-4C-alkoxy groups, which is substituted by a further 1-4C-alkoxy group. Examples which may be mentioned are the groups 2-(methoxy)ethoxy ($\text{CH}_3\text{-O-CH}_2\text{-CH}_2\text{-O-}$) and 2-(ethoxy)ethoxy ($\text{CH}_3\text{-CH}_2\text{-O-CH}_2\text{-CH}_2\text{-O-}$).

1-4C-Alkoxy-1-4C-alkoxy-1-4C-alkyl represents one of the aforementioned 1-4C-alkoxy-1-4C-alkyl groups, which is substituted by one of the aforementioned 1-4C-alkoxy groups. An example which may be mentioned is the group 2-(methoxy)ethoxymethyl ($\text{CH}_3\text{-O-CH}_2\text{-CH}_2\text{-O-CH}_2\text{-}$).

Fluoro-1-4C-alkoxy represents one of the aforementioned 1-4C-alkoxy groups, which is completely or mainly substituted by fluorine, "mainly" meaning in this connection that more than half of the hydrogen atoms are replaced by fluorine atoms. Examples of completely or mainly fluoro-substituted 1-4C-alkoxy groups which may be mentioned are the 1,1,1,3,3,3-hexafluoro-2-propoxy, the 2-trifluoromethyl-2-propoxy, the 1,1,1-trifluoro-2-propoxy, the perfluoro-tert-butoxy, the 2,2,3,3,4,4,4-heptafluoro-1-butoxy, the 4,4,4-trifluoro-1-butoxy, the 2,2,3,3,3-pentafluoropropoxy, the perfluoroethoxy, the 1,2,2-trifluoroethoxy, in particular the 1,1,2,2-tetrafluoroethoxy, the 2,2,2-trifluoroethoxy, the trifluoromethoxy and preferably the difluoromethoxy group.

Fluoro-1-4C-alkoxy-1-4C-alkyl represents one of the aforementioned 1-4C-alkyl groups, which is substituted by a fluoro-1-4C-alkoxy group. Examples of fluoro-1-4C-alkoxy-1-4C-alkyl groups are the 1,1,2,2-tetrafluoroethoxymethyl, the 2,2,2-trifluoroethoxymethyl, the trifluoromethoxyethyl and the difluoromethoxyethyl group.

1-7C-Alkyl represents straight-chain or branched alkyl groups having 1 to 7 carbon atoms. Examples which may be mentioned are the heptyl, isoheptyl (5-methylhexyl), hexyl, isohexyl (4-methylpentyl), neohexyl (3,3-dimethylbutyl), pentyl, isopentyl (3-methylbutyl), neopentyl (2,2-dimethylpropyl), butyl, isobutyl, sec-butyl, tert-butyl, propyl, isopropyl, ethyl and the methyl group.

Groups Ar which may be mentioned are, for example, the following substituents: 4-acetoxyphenyl, 4-acetamidophenyl, 2-methoxyphenyl, 3-methoxyphenyl, 4-methoxyphenyl, 3-benzyloxyphenyl, 4-benzyloxyphenyl, 3-benzyloxy-4-methoxyphenyl, 4-benzyloxy-3-methoxyphenyl, 3,5-bis(trifluoromethyl)phenyl, 4-butoxyphenyl, 2-chlorophenyl, 3-chlorophenyl, 4-chlorophenyl, 2-chloro-6-fluorophenyl, 3-chloro-4-fluorophenyl, 2-chloro-5-nitrophenyl, 4-chloro-3-nitrophenyl, 3-(4-chlorophenoxy)phenyl, 2,4-dichlorophenyl, 3,4-difluorophenyl, 2,4-dihydroxyphenyl, 2,6-dimethoxyphenyl, 3,4-dimethoxy-5-hydroxyphenyl, 2,5-dimethylphenyl, 3-ethoxy-4-hydroxyphenyl, 2-fluorophenyl, 4-fluorophenyl, 4-hydroxyphenyl, 2-hydroxy-5-nitrophenyl, 3-methoxy-2-nitrophenyl, 3-nitrophenyl, 2,3,5-trichlorophenyl, 2,4,6-trihydroxyphenyl, 2,3,4-trimethoxyphenyl, 2-hydroxy-1-naphthyl, 2-methoxy-1-naphthyl, 4-methoxy-1-naphthyl, 1-methyl-2-pyrrolyl, 2-pyrrolyl, 3-methyl-2-pyrrolyl, 3,4-dimethyl-2-pyrrolyl, 4-(2-methoxycarbonyl)ethyl-3-methyl-2-pyrrolyl, 5-ethoxycarbonyl-2,4-dimethyl-3-pyrrolyl, 3,4-dibromo-5-methyl-2-pyrrolyl, 2,5-dimethyl-1-phenyl-3-pyrrolyl, 5-carboxy-3-ethyl-4-methyl-2-pyrrolyl, 3,5-dimethyl-2-pyrrolyl, 2,5-dimethyl-1-(4-trifluoromethylphenyl)-3-pyrrolyl, 1-(2,6-dichloro-4-trifluoromethylphenyl)-2-pyrrolyl, 1-(2-nitrobenzyl)-2-

pyrrolyl, 1-(2-fluorophenyl)-2-pyrrolyl, 1-(4-trifluoromethoxyphenyl)-2-pyrrolyl, 1-(2-nitrobenzyl)-2-pyrrolyl, 1-(4-ethoxycarbonyl)-2,5-dimethyl-3-pyrrolyl, 5-chloro-1,3-dimethyl-4-pyrazolyl, 5-chloro-1-methyl-3-trifluoromethyl-4-pyrazolyl, 1-(4-chlorobenzyl)-5-pyrazolyl, 1,3-dimethyl-5-(4-chlorophenoxy)-4-pyrazolyl, 1-methyl-3-trifluoromethyl-5-(3-trifluoromethylphenoxy)-4-pyrazolyl, 4-methoxycarbonyl-1-(2,6-dichlorophenyl)-5-pyrazolyl, 5-allyloxy-1-methyl-3-trifluoromethyl-4-pyrazolyl, 5-chloro-1-phenyl-3-trifluoromethyl-4-pyrazolyl, 3,5-dimethyl-1-phenyl-4-imidazolyl, 4-bromo-1-methyl-5-imidazolyl, 2-butylimidazolyl, 1-phenyl-1,2,3-triazol-4-yl, 3-indolyl, 4-indolyl, 7-indolyl, 5-methoxy-3-indolyl, 5-benzyloxy-3-indolyl, 1-benzyl-3-indolyl, 2-(4-chlorophenyl)-3-indolyl, 7-benzyloxy-3-indolyl, 6-benzyloxy-3-indolyl, 2-methyl-5-nitro-3-indolyl, 4,5,6,7-tetrafluoro-3-indolyl, 1-(3,5-difluorobenzyl)-3-indolyl, 1-methyl-2-(4-trifluorophenoxy)-3-indolyl, 1-methyl-2-benzimidazolyl, 5-nitro-2-furyl, 5-hydroxymethyl-2-furyl, 2-furyl, 3-furyl, 5-(2-nitro-4-trifluoromethylphenyl)-2-furyl, 4-ethoxycarbonyl-5-methyl-2-furyl, 5-(2-trifluoromethoxyphenyl)-2-furyl, 5-(4-methoxy-2-nitrophenyl)-2-furyl, 4-bromo-2-furyl, 5-dimethylamino-2-furyl, 5-bromo-2-furyl, 5-sulfo-2-furyl, 2-benzofuryl, 2-thienyl, 3-thienyl, 3-methyl-2-thienyl, 4-bromo-2-thienyl, 5-bromo-2-thienyl, 5-nitro-2-thienyl, 5-methyl-2-thienyl, 5-(4-methoxyphenyl)-2-thienyl, 4-methyl-2-thienyl, 3-phenoxy-2-thienyl, 5-carboxy-2-thienyl, 2,5-dichloro-3-thienyl, 3-methoxy-2-thienyl, 2-benzothienyl, 3-methyl-2-benzothienyl, 2-bromo-5-chloro-3-benzothienyl, 2-thiazolyl, 2-amino-4-chloro-5-thiazolyl, 2,4-dichloro-5-thiazolyl, 2-diethylamino-5-thiazolyl, 3-methyl-4-nitro-5-isoxazolyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 6-methyl-2-pyridyl, 3-hydroxy-5-hydroxymethyl-2-methyl-4-pyridyl, 2,6-dichloro-4-pyridyl, 3-chloro-5-trifluoromethyl-2-pyridyl, 4,6-dimethyl-2-pyridyl, 4-(4-chlorophenyl)-3-pyridyl, 2-chloro-5-methoxycarbonyl-6-methyl-4-phenyl-3-pyridyl, 2-chloro-3-pyridyl, 6-(3-trifluoromethylphenoxy)-3-pyridyl, 2-(4-chlorophenoxy)-3-pyridyl, 2,4-dimethoxy-5-pyrimidinyl, 2-quinolinyl, 3-quinolinyl, 4-quinolinyl, 2-chloro-3-quinolinyl, 2-chloro-6-methoxy-3-quinolinyl, 8-hydroxy-2-quinolinyl and 4-isoquinolinyl.

2-4C-Alkenyloxy represents groups, which in addition to the oxygen atom contain one of the abovementioned 2-4C-alkenyl groups. Examples, which may be mentioned, are the 2-butenyloxy, 3-butenyloxy, 1-propenyloxy and the 2-propenyloxy group (allyloxy group).

1-4C-Alkylcarbonyl represents a group, which in addition to the carbonyl group contains one of the aforementioned 1-4C-alkyl groups. An example which may be mentioned is the acetyl group.

Carboxy-1-4C-alkyl represents 1-4C-alkyl groups which are substituted by a carboxyl group. Examples, which may be mentioned, are the carboxymethyl and the 2-carboxyethyl group.

1-4C-Alkoxycarbonyl-1-4C-alkyl represents 1-4C-alkyl groups, which are substituted by one of the abovementioned 1-4C-alkoxycarbonyl groups. Examples, which may be mentioned, are the Methoxycarbonylmethyl and the ethoxycarbonylmethyl group.

Aryl-1-4C-alkyl represents one of the aforementioned 1-4C-alkyl groups, which is substituted by one of the abovementioned aryl groups. An exemplary preferred aryl-1-4C-alkyl group is the benzyl group.

Aryl-1-4C-alkoxy represents one of the aforementioned 1-4C-alkoxy groups, which is substituted by one of the abovementioned aryl groups. An exemplary preferred aryl-1-4C-alkoxy group is the benzyloxy group.

1-4C-Alkylcarbonylamino represents an amino group to which a 1-4C-alkylcarbonyl group is bonded. Examples which may be mentioned are the propionylamino ($\text{C}_3\text{H}_7\text{C(O)NH-}$) and the acetyl amino group (acetamido group) ($\text{CH}_3\text{C(O)NH-}$).

1-4C-Alkoxy carbonylamino represents an amino group, which is substituted by one of the aforementioned 1-4C-alkoxy carbonyl groups. Examples, which may be mentioned, are the ethoxycarbonylamino and the methoxycarbonylamino group.

1-4C-Alkoxy-1-4C-alkoxy carbonyl represents a carbonyl group, to which one of the aforementioned 1-4C-alkoxy-1-4C-alkoxy groups is bonded. Examples which may be mentioned are the 2-(methoxy)ethoxycarbonyl ($\text{CH}_3\text{-O-CH}_2\text{CH}_2\text{-O-CO-}$) and the 2-(ethoxy)ethoxycarbonyl group ($\text{CH}_3\text{CH}_2\text{-O-CH}_2\text{CH}_2\text{-O-CO-}$).


1-4C-Alkoxy-1-4C-alkoxy carbonylamino represents an amino group, which is substituted by one of the aforementioned 1-4C-alkoxy-1-4C-alkoxy carbonyl groups. Examples which may be mentioned are the 2-(methoxy)ethoxycarbonylamino and the 2-(ethoxy)ethoxycarbonylamino group.

Possible salts of compounds of the formula 1 - depending on substitution - are especially all acid addition salts. Particular mention may be made of the pharmacologically tolerable salts of the inorganic and organic acids customarily used in pharmacy. Those suitable are water-soluble and water-insoluble acid addition salts with acids such as, for example, hydrochloric acid, hydrobromic acid, phosphoric acid, nitric acid, sulfuric acid, acetic acid, citric acid, D-gluconic acid, benzoic acid, 2-(4-hydroxybenzoyl)benzoic acid, butyric acid, sulfosalicylic acid, maleic acid, lauric acid, malic acid, fumaric acid, succinic acid, oxalic acid, tartaric acid, embonic acid, stearic acid, toluenesulfonic acid, methanesulfonic acid or 3-hydroxy-2-naphthoic acid, where the acids are used in salt preparation - depending on whether a mono- or polybasic acid is concerned and on which salt is desired - in an equimolar quantitative ratio or one differing there from.

Pharmacologically intolerable salts, which can initially be obtained, for example, as process products in the production of the compounds according to the invention on the industrial scale, are converted into the pharmacologically tolerable salts by processes known to the person skilled in the art.


It is known to the person skilled in the art that the compounds according to invention and their salts, if, for example, they are isolated in crystalline form, can contain various amounts of solvents. The invention therefore also comprises all solvates and in particular all hydrates of the compounds of the formula 1, and also all solvates and in particular all hydrates of the salts of the compounds of the formula 1.

One embodiment of the invention (embodiment a) relates to compounds of the formula 1a



in which R1, R2, R3 and Ar have the meanings given above, and their salts.

Another embodiment of the invention (embodiment b) relates to compounds of the formula 1b



in which R1, R2, R3 and Ar have the meanings given above, and their salts.

Among the compounds of formula 1, those are to be mentioned particularly, in which

R1 is hydrogen or 1-4C-alkyl,

R2 is hydrogen or 1-4C-alkyl,

R3 is halogen, carboxyl, 1-4C-alkoxycarbonyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy-1-4C-alkyl, or the group -CO-NR31R32,

where

R31 is hydrogen, 1-7C-alkyl, hydroxy-1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl and

R32 is hydrogen or 1-7C-alkyl,

or where

R31 and R32 together, including the nitrogen atom to which both are bonded, are a pyrrolidino, piperidino, piperazino, N-1-4C-alkylpiperazino or morpholino group,

X is O (oxygen) or NH,

Ar is a phenyl group substituted in the 2-position by R4 and in the 6-position by R5,
where


R4 is hydrogen, 1-4C-alkyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy, 1-4C-alkoxycarbonyl, trifluoromethyl, amino, mono- or di-1-4C-alkylamino, 1-4C-alkylcarbonylamino, 1-4C-alkoxycarbonylamino or 1-4C-alkoxy-1-4C-alkoxycarbonylamino and
R5 is hydrogen, 1-4C-alkyl or 1-4C-alkoxy,

or

Ar is selected from the group consisting of 4-acetoxypheyl, 4-acetamidophenyl, 2-methoxyphenyl, 3-methoxyphenyl, 4-methoxyphenyl, 3-benzyloxyphenyl, 4-benzyloxyphenyl, 3-benzyloxy-4-methoxyphenyl, 4-benzyloxy-3-methoxyphenyl, 3,5-bis(trifluoromethyl)phenyl, 4-butoxyphenyl, 2-chlorophenyl, 3-chlorophenyl, 4-chlorophenyl, 2-chloro-6-fluorophenyl, 3-chloro-4-fluorophenyl, 2-chloro-5-nitrophenyl, 4-chloro-3-nitrophenyl, 3-(4-chlorophenoxy)phenyl, 2,4-dichlorophenyl, 3,4-difluorophenyl, 2,4-dihydroxyphenyl, 2,6-dimethoxyphenyl, 3,4-dimethoxy-5-hydroxyphenyl, 2,5-dimethylphenyl, 3-ethoxy-4-hydroxyphenyl, 2-fluorophenyl, 4-fluorophenyl, 4-hydroxyphenyl, 2-hydroxy-5-nitrophenyl, 3-methoxy-2-nitrophenyl, 3-nitrophenyl, 2,3,5-trichlorophenyl, 2,4,6-trihydroxyphenyl, 2,3,4-trimethoxyphenyl, 2-hydroxy-1-naphthyl, 2-methoxy-1-naphthyl, 4-methoxy-1-naphthyl, 1-methyl-2-pyrrolyl, 2-pyrrolyl, 3-methyl-2-pyrrolyl, 3,4-dimethyl-2-pyrrolyl, 4-(2-methoxycarbonyl)ethyl-3-methyl-2-pyrrolyl, 5-ethoxycarbonyl-2,4-dimethyl-3-pyrrolyl, 3,4-dibromo-5-methyl-2-pyrrolyl, 2,5-dimethyl-1-phenyl-3-pyrrolyl, 5-carboxy-3-ethyl-4-methyl-2-pyrrolyl, 3,5-dimethyl-2-pyrrolyl, 2,5-dimethyl-1-(4-trifluoromethylphenyl)-3-pyrrolyl, 1-(2,6-dichloro-4-trifluoromethylphenyl)-2-pyrrolyl, 1-(2-nitrobenzyl)-2-pyrrolyl, 1-(2-fluorophenyl)-2-pyrrolyl, 1-(4-trifluoromethoxyphenyl)-2-pyrrolyl, 1-(2-nitrobenzyl)-2-pyrrolyl, 1-(4-ethoxycarbonyl)-2,5-dimethyl-3-pyrrolyl, 5-chloro-1,3-dimethyl-4-pyrazolyl, 5-chloro-1-methyl-3-trifluoromethyl-4-pyrazolyl, 1-(4-chlorobenzyl)-5-pyrazolyl, 1,3-dimethyl-5-(4-chlorophenoxy)-4-pyrazolyl, 1-methyl-3-trifluoromethyl-5-(3-trifluoromethylphenoxy)-4-pyrazolyl, 4-methoxycarbonyl-1-(2,6-dichlorophenyl)-5-pyrazolyl, 5-allyloxy-1-methyl-3-trifluoromethyl-4-pyrazolyl, 5-chloro-1-phenyl-3-trifluoromethyl-4-pyrazolyl, 3,5-dimethyl-1-phenyl-4-imidazolyl, 4-bromo-1-methyl-5-imidazolyl, 2-butylimidazolyl, 1-phenyl-1,2,3-triazol-4-yl, 3-indolyl, 4-indolyl, 7-indolyl, 5-methoxy-3-indolyl, 5-benzyloxy-3-indolyl, 1-benzyl-3-indolyl, 2-(4-chlorophenyl)-3-indolyl, 7-benzyloxy-3-indolyl, 6-benzyloxy-3-indolyl, 2-methyl-5-nitro-3-indolyl, 4,5,6,7-tetrafluoro-3-indolyl, 1-(3,5-difluorobenzyl)-3-indolyl, 1-methyl-2-(4-trifluorophenoxy)-3-indolyl, 1-methyl-2-benzimidazolyl, 5-nitro-2-furyl, 5-hydroxymethyl-2-furyl, 2-furyl, 3-furyl, 5-(2-nitro-4-trifluoromethylphenyl)-2-furyl, 4-ethoxycarbonyl-5-methyl-2-furyl, 5-(2-trifluoromethoxyphenyl)-2-furyl, 5-(4-methoxy-2-nitrophenyl)-2-furyl, 4-bromo-2-furyl, 5-dimethylamino-2-furyl, 5-bromo-2-furyl, 5-sulfo-2-furyl, 2-benzofuryl, 2-thienyl, 3-thienyl, 3-methyl-2-thienyl, 4-bromo-2-thienyl, 5-bromo-2-thienyl, 5-nitro-2-thienyl, 5-methyl-2-thienyl, 5-(4-methoxyphenyl)-2-thienyl, 4-methyl-2-thienyl, 3-phenoxy-2-thienyl, 5-carboxy-2-thienyl, 2,5-dichloro-3-thienyl, 3-methoxy-2-thienyl, 2-benzothienyl, 3-methyl-2-benzothienyl, 2-bromo-5-chloro-3-benzothienyl, 2-thiazolyl, 2-amino-4-chloro-5-thiazolyl, 2,4-dichloro-5-thiazolyl, 2-diethylamino-5-thiazolyl, 3-methyl-4-nitro-5-isoxazolyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 6-methyl-2-pyridyl, 3-hydroxy-5-hydroxymethyl-2-methyl-4-pyridyl, 2,6-dichloro-4-pyridyl, 3-chloro-5-trifluoromethyl-2-pyridyl, 4,6-dimethyl-2-pyridyl, 4-(4-chlorophenyl)-3-pyridyl, 2-chloro-5-methoxycarbonyl-6-methyl-4-phenyl-3-pyridyl,

2-chloro-3-pyridyl, 6-(3-trifluoromethylphenoxy)-3-pyridyl, 2-(4-chlorophenoxy)-3-pyridyl, 2,4-dimethoxy-5-pyrimidinyl, 2-quinolinyl, 3-quinolinyl, 4-quinolinyl, 2-chloro-3-quinolinyl, 2-chloro-6-methoxy-3-quinolinyl, 8-hydroxy-2-quinolinyl and 4-isoquinolinyl, and the salts of these compounds.

Among the compounds of the formula 1, those of the formula 1-1 have to be highlighted



in which

R1 is hydrogen or 1-4C-alkyl,

R2 is hydrogen or 1-4C-alkyl,

R3 is halogen, carboxyl, 1-4C-alkoxycarbonyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy-1-4C-alkyl or the group -CO-NR₃₁R₃₂,

where

R₃₁ is hydrogen, 1-7C-alkyl, hydroxy-1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl and

R₃₂ is hydrogen or 1-7C-alkyl,

or where

R₃₁ and R₃₂ together, including the nitrogen atom to which both are bonded, are a pyrrolidino, piperidino, piperazino, N-1-4C-alkylpiperazino or morpholino group,

R4 is hydrogen, 1-4C-alkyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy, 1-4C-alkoxycarbonyl, trifluoromethyl, amino, mono- or di-1-4C-alkylamino, 1-4C-alkylcarbonylamino, 1-4C-alkoxycarbonylamino or 1-4C-alkoxy-1-4C-alkoxycarbonylamino,

R5 is hydrogen, 1-4C-alkyl or 1-4C-alkoxy and

X is O (oxygen) or NH,

and the salts of these compounds.

Compounds of embodiment a to be highlighted are those of formula 1-1, in which

R1 is hydrogen or 1-4C-alkyl,

R2 is hydrogen or 1-4C-alkyl,

- R3 is halogen, carboxyl, 1-4C-alkoxycarbonyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy-1-4C-alkyl or the group -CO-NR₃₁R₃₂,
 where
 R₃₁ is hydrogen, 1-7C-alkyl, hydroxy-1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl and
 R₃₂ is hydrogen or 1-7C-alkyl,
 or where
 R₃₁ and R₃₂ together, including the nitrogen atom to which both are bonded, are a pyrrolidino, piperidino, piperazino, N-1-4C-alkylpiperazino or morpholino group,
 R4 is hydrogen, 1-4C-alkyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy, 1-4C-alkoxycarbonyl, trifluoromethyl, amino, mono- or di-1-4C-alkylamino, 1-4C-alkylcarbonylamino, 1-4C-alkoxycarbonylamino or 1-4C-alkoxy-1-4C-alkoxycarbonylamino,
 R5 is hydrogen, 1-4C-alkyl or 1-4C-alkoxy and
 X is O (oxygen),
 and the salts of these compounds.

Compounds of embodiment b to be highlighted are those of formula 1-1, in which

- R1 is hydrogen or 1-4C-alkyl,
 R2 is hydrogen or 1-4C-alkyl,
 R3 is halogen, carboxyl, 1-4C-alkoxycarbonyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy-1-4C-alkyl or the group -CO-NR₃₁R₃₂,
 where
 R₃₁ is hydrogen, 1-7C-alkyl, hydroxy-1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl and
 R₃₂ is hydrogen or 1-7C-alkyl,
 or where
 R₃₁ and R₃₂ together, including the nitrogen atom to which both are bonded, are a pyrrolidino, piperidino, piperazino, N-1-4C-alkylpiperazino or morpholino group,
 R4 is hydrogen, 1-4C-alkyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy, 1-4C-alkoxycarbonyl, trifluoromethyl, amino, mono- or di-1-4C-alkylamino, 1-4C-alkylcarbonylamino, 1-4C-alkoxycarbonylamino or 1-4C-alkoxy-1-4C-alkoxycarbonylamino,
 R5 is hydrogen, 1-4C-alkyl or 1-4C-alkoxy and
 X is NH,
 and the salts of these compounds.

Compounds of the formula 1-1 to be emphasised are those, in which

- R1 is 1-4C-alkyl,
 R2 is 1-4C-alkyl,
 R3 is carboxyl, 1-4C-alkoxycarbonyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy-1-4C-alkyl or the group -CO-NR₃₁R₃₂,
 where
 R₃₁ is hydrogen, 1-7C-alkyl, hydroxy-1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl and
 R₃₂ is hydrogen or 1-7C-alkyl,

or where

R31 and R32 together, including the nitrogen atom to which both are bonded, are a pyrrolidino, piperidino, piperazino, N-1-4C-alkylpiperazino or morpholino group,

R4 is hydrogen, 1-4C-alkyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy, 1-4C-alkoxycarbonyl, trifluoromethyl, amino, mono- or di-1-4C-alkylamino, 1-4C-alkylcarbonylamino, 1-4C-alkoxycarbonylamino or 1-4C-alkoxy-1-4C-alkoxycarbonylamino,

R5 is hydrogen, 1-4C-alkyl or 1-4C-alkoxy and

X is O (oxygen) or NH,

and the salts of these compounds.

Preferred compounds of the formula 1-1 are those, in which

R1 is 1-4C-alkyl,

R2 is 1-4C-alkyl,

R3 is carboxyl, 1-4C-alkoxycarbonyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy-1-4C-alkyl or the group -CO-NR31R32,

where

R31 is hydrogen, 1-4C-alkyl, hydroxy-1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl and

R32 is hydrogen or 1-4C-alkyl,

R4 is 1-4C-alkyl or 1-4C-alkylcarbonylamino,

R5 is 1-4C-alkyl and

X is O (oxygen) or NH,

and their salts.

Particularly preferred compounds of the formula 1-1 are those, in which

R1 is 1-4C-alkyl,

R2 is 1-4C-alkyl,

R3 is carboxyl, 1-4C-alkoxycarbonyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy-1-4C-alkyl or the group -CO-NR31R32,

where

R31 is hydrogen, 1-4C-alkyl or hydroxy-1-4C-alkyl and

R32 is hydrogen or 1-4C-alkyl,

R4 is 1-4C-alkyl,

R5 is 1-4C-alkyl and

X is O (oxygen) or NH,


and their salts.

Particularly preferred are the compounds given as final products of formula 1 in the examples, and the salts of these compounds.

The compounds according to the invention can be synthesised from corresponding starting compounds, for example according to the reaction schemes given below. The synthesis is carried out in a manner known to the expert, for example as described in more detail in the following examples.


According to the invention, the compounds of formula 1 can be prepared as outlined in the reaction schemes 1 and 2, which illustrate processes known to the expert and which use known starting materials. The particular method for the synthesis and reaction sequence of the compounds of formula 1 is chosen having regard to the specific nature of the substituents and their position. One of the processes for producing the compounds of formula 1 consists in condensing a 3,5-disubstituted 2-aminopyrazine II with an alpha-halocarbonyl compound III (scheme 1). The required 3,5-disubstituted 2-aminopyrazines II are obtained by a substitution reaction of a 5-substituted 2-amino-3-bromopyrazine I, containing any desired substituent R₃, with ArCH₂XH (X = O or NH), analogously to known procedures (see, for example, EP 204285).

Scheme 1.



Another process (scheme 2) for producing the compounds of formula 1 consists in carrying out a substitution reaction starting with an appropriate substituted imidazo[1,2-a]pyrazine (IV or V). It is thus possible, for example, starting from compounds of formula IV, to prepare compounds of formula 1 where R₂ = CH₂OH (e.g. by Vilsmeier reaction and subsequent reduction) or where R₂ = Br or Cl (by bromination or chlorination). Further derivatization of compounds of formula 1, where R₂ = Br or Cl, can be accomplished for example by metal-catalysed carbonylation to get compounds of formula 1 where R₂ = alkoxycarbonyl or by using the Sonogashira reaction to get compounds of formula 1 where R₂ = propynyl. A substitution of R₃ can be carried out likewise, for example by palladium-catalysed carbonylation of compounds V as described in more detail in the examples. If compounds, where R₃ = -CO-NR₃₁R₃₂, are desired, an appropriate derivatization can be performed in a manner known per se (conversion of an ester or carboxylic acid into an amide).

Scheme 2.



The following examples serve to illustrate the invention in greater detail without restricting it. Likewise, further compounds of the formula 1 whose preparation is not described explicitly can be prepared in an analogous manner or in a manner familiar per se to the person skilled in the art using customary process techniques. The abbreviation min stands for minute(s), h for hour(s). The compounds under the scope of formula 1 named expressly as examples, and any salts of these compounds, are preferred subject matter of the invention.

Examples

1. 2-Ethyl-6-methyl-benzylamine

To a suspension of 10.4 g (275 mmol) of lithium aluminium hydride in 200 ml of dried diethyl ether is slowly added a solution of 20.0 g (138 mmol) of 2-ethyl-6-methyl-benzonitrile in 60 ml of diethyl ether at -10°C . After 1 h at 0°C and 1 h at room temperature, the reaction mixture is carefully hydrolyzed with 4 ml of water and 4 ml of 6N sodium hydroxide solution. After 2 h at room temperature, anhydrous magnesium sulfate is added and the reaction mixture is filtered through Celite. Evaporation of the solvent yields 15.5 g (80 %) of the title compound as a colourless oil which is used without further purification in the next step.

2. 2-Amino-5-bromo-3-(2-ethyl-6-methyl-benzylamino)-pyrazine

A solution of 1.26 g (5 mmol) of 2-amino-3,5-dibromopyrazine (B. Jiang et al., *Bioorg. Med. Chem.* **2001**, *9*, 1149-1154), 1.5 g (10 mmol) of 2-ethyl-6-methyl-benzylamine and 1.5 ml of triethylamine in 3.5 ml of acetonitrile in a sealed tube is irradiated in a microwave-oven for 40 min (temperature 180°C). The crude reaction mixtures of 10 such runs are combined, treated with saturated sodium hydrogen carbonate and extracted with ethyl acetate. The organic phase is dried over anhydrous magnesium sulfate and evaporated. The residue is purified by column chromatography on silica gel using light petroleum ether:ethyl acetate (4:1, v/v). Crystallization from dioxane yields 10.2 g (68 %) of the title compound as a colourless solid (m.p. 155°C).

3. 6-Bromo-8-(2-ethyl-6-methyl-benzylamino)-2,3-dimethyl-imidazo[1,2-a]pyrazine oxalate

To a suspension of 10.0 g (31 mmol) of 2-amino-5-bromo-3-(2-ethyl-6-methyl-benzylamino)-pyrazine in 60 ml of dioxane are added 4.9 ml (46.7 mmol) of 3-bromo-2-butanone and the resulting mixture is heated to 100°C . After 2 h a further amount of 4.9 ml (46.7 mmol) of 3-bromo-2-butanone is added and the mixture is stirred for 16 h. The mixture is cooled down, diluted with dichloromethane and extracted with saturated aqueous sodium hydrogen carbonate. The organic phase is dried over anhydrous magnesium sulfate and evaporated. Purification of the residue by column chromatography on silica gel using light petroleum ether/ethyl acetate (4:1, v/v) gives a colourless oil which is dissolved in acetone and treated with a solution of 3.91 g (31 mmol) of oxalic acid dihydrate in acetone. The precipitate is collected and washed with n-heptane to yield 10 g (70 %) of the title compound as a colourless solid (m.p. 163°C).

4. Ethyl 8-(2-ethyl-6-methyl-benzylamino)-2,3-dimethyl-imidazo[1,2-a]pyrazine-6-carboxylate

10.0 g (22 mmol) of 6-bromo-8-(2-ethyl-6-methyl-benzylamino)-2,3-dimethyl-imidazo[1,2-a]pyrazine oxalate are treated with saturated aqueous sodium hydrogen carbonate and extracted with ethyl acetate. The organic phase is separated, dried over anhydrous magnesium sulfate and evaporated to give 6-bromo-8-(2-ethyl-6-methyl-benzylamino)-2,3-dimethyl-imidazo[1,2-a]pyrazine as a colourless oil. The resulting oil thus obtained is dissolved in 80 ml of absolute ethanol and 16 ml of triethylamine and transferred to an autoclave. After addition of 0.5 g (2.2 mmol) of palladium(II) acetate and 1.64 g

(6.2 mmol) of triphenylphosphine, the reaction mixture is carbonylated (10 bar carbon monoxide pressure, 100 °C) for 14 h. The reaction mixture is cooled down, filtered and evaporated to leave an orange coloured oil which is dissolved in dichloromethane and extracted with water. The organic phase is dried over anhydrous magnesium sulfate and evaporated. Purification of the residue by crystallization from ethyl acetate/n-heptane yields 7.2 g (89 %) of the title compound as a colourless solid (m.p. 144 °C).

5. 8-(Dimethylaminocarbonyl)-8-(2-ethyl-6-methyl-benzylamino)- 2,3-dimethyl-imidazo[1,2-a]-pyrazine

To a solution of 2.3 g (5.1 mmol) of 6-bromo-8-(2-ethyl-6-methyl-benzylamino)- 2,3-dimethyl-imidazo[1,2-a]pyrazine in 50 ml of dimethylamine (2M solution in tetrahydrofuran) are added 0.17 g (0.76 mmol) of palladium(II) acetate and 0.8 g (3.1 mmol) of triphenylphosphine. The mixture is transferred to an autoclave and carbonylated (6 bar carbon monoxide pressure, 120 °C) for 16 h. The reaction mixture is cooled down, evaporated and the residue dissolved in dichloromethane. The organic phase is extracted with saturated aqueous ammonium chloride solution, dried over anhydrous magnesium sulfate and evaporated. Purification of the residue by column chromatography on silica gel using light petroleum ether/ethyl acetate (1:1, v/v) yields 1.22 g (66 %) of the title compound as a colourless solid (m.p. 174 °C).

6. 8-(2-Ethyl-6-methyl-benzylamino)- 2,3-dimethyl-imidazo[1,2-a]pyrazine-6-carboxylic acid

To a solution of 4.0 g (10.9 mmol) of ethyl 8-(2-ethyl-6-methyl-benzylamino)- 2,3-dimethyl-imidazo[1,2-a]pyrazine-6-carboxylate in 40 ml of dioxane are added 8 ml of 2N aqueous sodium hydroxide solution. After 1 h at 80 °C, the reaction mixture is evaporated to half of its volume and the pH is adjusted to 6 by the addition of 6N hydrochloric acid. The thick precipitate is collected, washed with water and dried *in vacuo* over phosphorus pentoxide to yield 3.52 g (95 %) of the title compound as a colourless solid (m.p. 230 °C).

7. 8-(2-Ethyl-6-methyl-benzylamino)-6-(pyrrolidinocarbonyl)- 2,3-dimethyl-imidazo[1,2-a]pyrazine

To a suspension of 0.5 g (1.48 mmol) of 8-(2-ethyl-6-methyl-benzylamino)- 2,3-dimethyl-imidazo[1,2-a]pyrazine-6-carboxylic acid in 10 ml of dichloromethane are added 0.7 g (2.2 mmol) of O-(1H-benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium tetrafluoroborate (TBTU). After 30 min 0.5 ml (6 mmol) of pyrrolidine are added and the mixture is stirred for 7 h. The reaction mixture is extracted with 2N aqueous sodium hydroxide solution, the organic phase is separated, dried over anhydrous magnesium sulfate and evaporated. Purification of the residue by column chromatography on silica gel using dichloromethane/methanol (20:1, v/v) and crystallization from ethyl acetate/n-heptane yields 0.45 g (78 %) of the title compound as a colourless solid (m.p. 197 °C).

8. 8-(2-Ethyl-6-methyl-benzylamino)- 2,3-dimethyl-imidazo[1,2-a]pyrazine-6-carboxamide

To a suspension of 1.02 g (3 mmol) of 8-(2-ethyl-6-methyl-benzylamino)-2,3-dimethyl-imidazo[1,2-a]pyrazine-6-carboxylic acid in 20 ml of dichloromethane are added 1.61 g (5 mmol) of O-(1H-benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium tetrafluoroborate (TBTU). After 30 min ammonia gas is passed over the mixture. After 1 h, a further amount of 1.0 g (3.1 mmol) of TBTU is added. Stirring is continued for 1 h at room temperature and finally 1 h under reflux. The reaction mixture is extracted with 2N aqueous sodium hydroxide solution, the organic phase is separated, dried over anhydrous magnesium sulfate and evaporated. Purification of the residue by column chromatography on silica gel using dichloromethane/methanol (20:1, v/v) and crystallization from ethyl acetate/n-heptane yields 0.67 g (66 %) of the title compound as a colourless solid (m.p. 227 °C).

9. 8-(2-Ethyl-6-methyl-benzylamino)-6-(methylaminocarbonyl)-2,3-dimethyl-imidazo[1,2-a]-pyrazine

To a suspension of 1.02 g (3 mmol) of 8-(2-ethyl-6-methyl-benzylamino)-2,3-dimethyl-imidazo[1,2-a]pyrazine-6-carboxylic acid in 20 ml of dichloromethane are added 1.61 g (5 mmol) of O-(1H-benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium tetrafluoroborate (TBTU). After 30 min stirring at room temperature 1 ml (8 mmol) of methylamine (8M in ethanol) is added. After 1 h, a further amount of 0.5 ml (4 mmol) of methylamine (8M in ethanol) is added and stirring is continued for 16 h. The reaction mixture is extracted with 2N aqueous sodium hydroxide solution, the organic phase is separated, dried over anhydrous magnesium sulfate and evaporated. Purification of the residue by column chromatography on silica gel using ethyl acetate/light petroleum ether (1:1, v/v) and crystallization from ethyl acetate/n-heptane yields 0.99 g (94 %) of the title compound as a colourless solid (m.p. 120 °C).

10. 8-(2-Ethyl-6-methyl-benzylamino)-6-(2-methoxyethylaminocarbonyl)-2,3-dimethyl-imidazo[1,2-a]pyrazine

A solution of 1.0 g (2.73 mmol) of ethyl 8-(2-ethyl-6-methyl-benzylamino)-2,3-dimethyl-imidazo[1,2-a]pyrazine-6-carboxylate in 10 ml of 2-methoxyethylamine is heated under reflux for 20 h. The reaction mixture is diluted with water and extracted with dichloromethane. The organic phase is dried over anhydrous magnesium sulfate and evaporated. Purification of the residue by column chromatography on silica gel using ethyl acetate/light petroleum ether (1:1, v/v) and crystallization from diisopropyl ether yields 0.53 g (49 %) of the title compound as a colourless solid (m.p. 111 °C).

11. 8-(2-Ethyl-6-methyl-benzylamino)-6-(2-hydroxyethylaminocarbonyl)-2,3-dimethyl-imidazo[1,2-a]pyrazine

A suspension of 1.1 g (3 mmol) of ethyl 8-(2-ethyl-6-methyl-benzylamino)-2,3-dimethyl-imidazo[1,2-a]pyrazine-6-carboxylate in 10 ml of 2-aminoethanol is heated to 80 °C for 30 min. The reaction mixture is diluted with an additional amount of 10 ml of 2-aminoethanol and the temperature is raised to 100 °C. After 1 h, the reaction mixture is cooled down and the precipitate is collected and washed with water. The colourless solid is dried *in vacuo* over phosphorus pentoxide to yield 1.04 g (91 %) of the title compound (m.p. 229 °C).

12. 8-(2-Ethyl-6-methyl-benzylamino)-6-(hydroxymethyl)-2,3-dimethyl-imidazo[1,2-a]pyrazine

To a suspension of 0.31 g (8.2 mmol) of lithium aluminium hydride in 10 ml of dried tetrahydrofuran is slowly added a solution of 1.0 g (2.7 mmol) of ethyl 8-(2-ethyl-6-methyl-benzylamino)-2,3-dimethyl-imidazo[1,2-a]pyrazine-6-carboxylate in 20 ml of tetrahydrofuran at 0 °C. After 1 h at 0 °C, the reaction mixture is carefully hydrolyzed with 0.2 ml of water, 0.4 ml of 6N sodium hydroxide solution and 1 ml of water. After 1 h at room temperature, anhydrous magnesium sulfate is added and the reaction mixture is filtered through Celite. On evaporation of the filtrate, a precipitate is obtained which is washed with diethyl ether and dried *in vacuo* to yield 0.77 g (87 %) of the title compound as a colourless solid (m.p. 166 °C).

13. 8-(2-Ethyl-6-methyl-benzylamino)-6-(methoxymethyl)-2,3-dimethyl-imidazo[1,2-a]pyrazine hydrochloride

To a suspension of 0.5 g (1.54 mmol) of 8-(2-ethyl-6-methyl-benzylamino)-6-hydroxymethyl-2,3-dimethyl-imidazo[1,2-a]pyrazine in 5 ml of absolute N,N-dimethylformamide are added 0.18 g (4.5 mmol) of sodium hydride (60 % w/w dispersion in mineral oil) in portions at room temperature. After 30 min, 0.12 ml (1.95 mmol) of methyl iodide are slowly added. After 30 min, the reaction mixture is carefully hydrolyzed with saturated aqueous sodium hydrogen carbonate solution and extracted with dichloromethane. The organic phase is dried over anhydrous magnesium sulfate and evaporated. Purification of the residue by column chromatography on silica gel using ethyl acetate/light petroleum ether (1:4, v/v) yields 0.18 g of a colourless oil which is dissolved in dichloromethane and treated with hydrogen chloride (1.5M in diethyl ether). Evaporation of all volatiles yields 0.12 g (21 %) of the title compound as a colourless solid (m.p. 177 °C).

Commercial utility

The compounds of the formula 1 and their salts have valuable pharmacological properties which make them commercially utilizable. In particular, they exhibit marked inhibition of gastric acid secretion and an excellent gastric and intestinal protective action in warm-blooded animals, in particular humans. In this connection, the compounds according to the invention are distinguished by a high selectivity of action, an advantageous duration of action, a particularly good enteral activity, the absence of significant side effects and a large therapeutic range.

"Gastric and intestinal protection" in this connection is understood as meaning the prevention and treatment of gastrointestinal diseases, in particular of gastrointestinal inflammatory diseases and lesions (such as, for example, gastric ulcer, peptic ulcer, including peptic ulcer bleeding, duodenal ulcer, gastritis, hyperacidic or medicament-related functional dyspepsia), which can be caused, for example, by microorganisms (e.g. *Helicobacter pylori*), bacterial toxins, medicaments (e.g. certain antiinflammatories and antirheumatics, such as NSAIDs and COX-inhibitors), chemicals (e.g. ethanol), gastric acid or stress situations. "Gastric and intestinal protection" is understood to include, according to general knowledge, gastroesophageal reflux disease (GERD), the symptoms of which include, but are not limited to, heartburn and/or acid regurgitation.

In their excellent properties, the compounds according to the invention surprisingly prove to be clearly superior to the compounds known from the prior art in various models in which the antiulcerogenic and the antisecretory properties are determined. On account of these properties, the compounds of the formula 1 and their pharmacologically acceptable salts are outstandingly suitable for use in human and veterinary medicine, where they are used, in particular, for the treatment and/or prophylaxis of disorders of the stomach and/or intestine.

A further subject of the invention are therefore the compounds according to the invention for use in the treatment and/or prophylaxis of the abovementioned diseases.

The invention likewise includes the use of the compounds according to the invention for the production of medicaments which are employed for the treatment and/or prophylaxis of the abovementioned diseases.

The invention furthermore includes the use of the compounds according to the invention for the treatment and/or prophylaxis of the abovementioned diseases.

A further subject of the invention are medicaments which comprise one or more compounds of the formula 1 and/or their pharmacologically acceptable salts.

The medicaments are prepared by processes which are known per se and familiar to the person skilled in the art. As medicaments, the pharmacologically active compounds according to the invention

(= active compounds) are either employed as such, or preferably in combination with suitable pharmaceutical auxiliaries or excipients in the form of tablets, coated tablets, capsules, suppositories, patches (e.g. as TTS), emulsions, suspensions or solutions, the active compound content advantageously being between 0.1 and 95% and it being possible to obtain a pharmaceutical administration form exactly adapted to the active compound and/or to the desired onset and/or duration of action (e.g. a sustained-release form or an enteric form) by means of the appropriate selection of the auxiliaries and excipients.

The auxiliaries and excipients which are suitable for the desired pharmaceutical formulations are known to the person skilled in the art on the basis of his/her expert knowledge. In addition to solvents, gel-forming agents, suppository bases, tablet auxiliaries and other active compound excipients, it is possible to use, for example, antioxidants, dispersants, emulsifiers, antifoams, flavor corrigents, preservatives, solubilizers, colorants or, in particular, permeation promoters and complexing agents (e.g. cyclodextrins).

The active compounds can be administered orally, parenterally or percutaneously.

In general, it has proven advantageous in human medicine to administer the active compound(s) in the case of oral administration in a daily dose of approximately 0.01 to approximately 20, preferably 0.05 to 5, in particular 0.1 to 1.5, mg/kg of body weight, if appropriate in the form of several, preferably 1 to 4, individual doses to achieve the desired result. In the case of a parenteral treatment, similar or (in particular in the case of the intravenous administration of the active compounds), as a rule, lower doses can be used. The establishment of the optimal dose and manner of administration of the active compounds necessary in each case can easily be carried out by any person skilled in the art on the basis of his/her expert knowledge.

If the compounds according to the invention and/or their salts are to be used for the treatment of the abovementioned diseases, the pharmaceutical preparations can also contain one or more pharmacologically active constituents of other groups of medicaments, for example: tranquilizers (for example from the group of the benzodiazepines, for example diazepam), spasmolytics (for example, biefamiverine or camylofine), anticholinergics (for example, oxyphenycyclimine or phencarbamide), local anesthetics, (for example, tetracaine or procaine), and, if appropriate, also enzymes, vitamins or amino acids.

To be emphasized in this connection is in particular the combination of the compounds according to the invention with pharmaceuticals which inhibit acid secretion, such as, for example, H₂ blockers (e.g. cimetidine, ranitidine), H⁺/K⁺ ATPase inhibitors (e.g. omeprazole, pantoprazole), or further with so-called peripheral anticholinergics (e.g. pirenzepine, telenzepine) and with gastrin antagonists with the aim of increasing the principal action in an additive or super-additive sense and/or of eliminating or of decreasing the side effects, or further the combination with antibacterially active substances (such as, for example, cephalosporins, tetracyclines, penicillins, macrolides, nitroimidazoles or alternatively

bismuth salts) for the control of *Helicobacter pylori*. Suitable antibacterial co-components which may be mentioned are, for example, mezlocillin, ampicillin, amoxicillin, cefalothin, cefoxitin, cefotaxime, imipenem, gentamycin, amikacin, erythromycin, ciprofloxacin, metronidazole, clarithromycin, azithromycin and combinations thereof (for example clarithromycin + metronidazole).

In view of their excellent gastric and intestinal protection action, the compounds of formula 1 are suited for a free or fixed combination with those medicaments (e.g. certain antiinflammatories and antirheumatics, such as NSAIDs), which are known to have a certain ulcerogenic potency. In addition, the compounds of formula 1 are suited for a free or fixed combination with motility-modifying drugs.

Pharmacology

The excellent gastric protective action and the gastric acid secretion-inhibiting action of the compounds according to the invention can be demonstrated in investigations on animal experimental models. The compounds according to the invention investigated in the model mentioned below have been provided with numbers which correspond to the numbers of these compounds in the examples.

Testing of the secretion-inhibiting action on the perfused rat stomach

In Table A which follows, the influence of the compounds according to the invention on the pentagastrin-stimulated acid secretion of the perfused rat stomach after intraduodenal administration in vivo is shown.

Table A

Example No.	Dose ($\mu\text{mol/kg}$) i.d.	Inhibition of acid secretion (%)
5	1	49
13	1	46

Methodology

The abdomen of anesthetized rats (CD rat, female, 200-250 g; 1.5 g/kg i.m. urethane) was opened after tracheotomy by a median upper abdominal incision and a PVC catheter was fixed transorally in the esophagus and another via the pylorus such that the ends of the tubes just projected into the gastric lumen. The catheter leading from the pylorus led outward into the right abdominal wall through a side opening.


After thorough rinsing (about 50-100 ml), warm (37°C) physiological NaCl solution was continuously passed through the stomach (0.5 ml/min, pH 6.8-6.9; Braun-Unita I). The pH (pH meter 632, glass electrode EA 147; $\phi = 5$ mm, Metrohm) and, by titration with a freshly prepared 0.01N NaOH solution to pH 7 (Dosimat 665 Metrohm), the secreted HCl were determined in the effluent in each case collected at an interval of 15 minutes.

The gastric secretion was stimulated by continuous infusion of 1 $\mu\text{g/kg}$ (= 1.65 ml/h) of i.v. pentagastrin (left femoral vein) about 30 min after the end of the operation (i.e. after determination of 2 preliminary fractions). The substances to be tested were administered intraduodenally in a 2.5 ml/kg liquid volume 60 min after the start of the continuous pentagastrin infusion.

The body temperature of the animals was kept at a constant 37.8-38°C by infrared irradiation and heat pads (automatic, stepless control by means of a rectal temperature sensor).

Patent claims

1. Compounds of the formula 1



in which

R1 is hydrogen, 1-4C-alkyl, 3-7C-cycloalkyl, 3-7C-cycloalkyl-1-4C-alkyl, 1-4C-alkoxy, 1-4C-alkoxy-1-4C-alkyl, 1-4C-alkoxycarbonyl, 2-4C-alkenyl, 2-4C-alkynyl, fluoro-1-4C-alkyl or hydroxy-1-4C-alkyl,

R2 is hydrogen, 1-4C-alkyl, aryl, 3-7C-cycloalkyl, 3-7C-cycloalkyl-1-4C-alkyl, 1-4C-alkoxycarbonyl, hydroxy-1-4C-alkyl, fluoro-1-4C-alkyl, halogen, 2-4C-alkenyl, 2-4C-alkynyl, amino, mono- or di-1-4C-alkylamino-1-4C-alkyl or cyanomethyl,

R3 is halogen, fluoro-1-4C-alkyl, 2-4C-alkenyl, 2-4C-alkynyl, carboxyl, cyano, 1-4C-alkoxycarbonyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy-1-4C-alkyl, 1-4C-alkoxy-1-4C-alkoxy-1-4C-alkyl, fluoro-1-4C-alkoxy-1-4C-alkyl or the group -CO-NR31R32,

where

R31 is hydrogen, hydroxyl, 1-7C-alkyl, hydroxy-1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl and

R32 is hydrogen, 1-7C-alkyl, hydroxy-1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl,

or where

R31 and R32 together, including the nitrogen atom to which both are bonded, are a pyrrolidino, piperidino, piperazino, N-1-4C-alkylpiperazino or morpholino group,

X is O (oxygen) or NH and

Ar is a mono- or bicyclic aromatic residue, substituted by R4, R5, R6 and R7, which is selected from the group consisting of phenyl, naphthyl, pyrrolyl, pyrazolyl, imidazolyl, 1,2,3-triazolyl, indolyl, benzimidazolyl, furyl, benzofuryl, thienyl, benzothienyl, thiazolyl, isoxazolyl, pyridinyl, pyrimidinyl, chinolinyl and isochinolinyl,

wherein

R4 is hydrogen, 1-4C-alkyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy, 2-4C-alkenyloxy, 1-4C-alkylcarbonyl, carboxy, 1-4C-alkoxycarbonyl, carboxy-1-4C-alkyl, 1-4C-alkoxycarbonyl-1-4C-alkyl, halogen, hydroxy, aryl, aryl-1-4C-alkyl, aryl-oxy, aryl-1-4C-alkoxy, trifluoromethyl, nitro, amino, mono- or di-1-4C-alkylamino, 1-4C-alkylcarbonylamino, 1-4C-alkoxycarbonylamino, 1-4C-alkoxy-1-4C-alkoxycarbonylamino or sulfonyl,

in which

aryl is phenyl or substituted phenyl with one, two or three identical or different substituents from the group of 1-4C-alkyl, 1-4C-alkoxy, carboxy, 1-4C-alkoxycarbonyl, halogen, trifluoromethyl, nitro, trifluoromethoxy, hydroxy and cyano,


R5 is hydrogen, 1-4C-alkyl, 1-4C-alkoxy, 1-4C-alkoxycarbonyl, halogen, trifluoromethyl or hydroxy,

R6 is hydrogen, 1-4C-alkyl, 1-4C-alkoxy, hydroxy or halogen and

R7 is hydrogen, 1-4C-alkyl or halogen,

and the salts of these compounds.

2. Compounds of formula 1 according to claim 1, in which X is O (oxygen).
3. Compounds of formula 1 according to claim 1, in which X is NH.
4. Compounds of the formula 1 according to claim 1, characterized by the formula 1-1



In which

R1 is hydrogen or 1-4C-alkyl,

R2 is hydrogen or 1-4C-alkyl,

R3 is halogen, carboxyl, 1-4C-alkoxycarbonyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy-1-4C-alkyl or the group -CO-NR31R32,

where

R31 is hydrogen, 1-7C-alkyl, hydroxy-1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl and

R32 is hydrogen or 1-7C-alkyl,

or where

R31 and R32 together, including the nitrogen atom to which both are bonded, are a pyrrolidino, piperidino, piperazino, N-1-4C-alkylpiperazino or morpholino group,

R4 is hydrogen, 1-4C-alkyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy, 1-4C-alkoxycarbonyl, trifluoromethyl, amino, mono- or di-1-4C-alkylamino, 1-4C-alkylcarbonylamino, 1-4C-alkoxycarbonylamino or 1-4C-alkoxy-1-4C-alkoxycarbonylamino,

R5 is hydrogen, 1-4C-alkyl or 1-4C-alkoxy and

X is O (oxygen) or NH,
and the salts of these compounds.

5. Compounds of the formula 1-1 according to claim 4, in which

R1 is 1-4C-alkyl,

R2 is 1-4C-alkyl,

R3 is carboxyl, 1-4C-alkoxycarbonyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy-1-4C-alkyl or the group -CO-NR31R32,

where

R31 is hydrogen, 1-7C-alkyl, hydroxy-1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl and

R32 is hydrogen or 1-7C-alkyl,

or where

R31 and R32 together, including the nitrogen atom to which both are bonded, are a pyrrolidino, piperidino, piperazino, N-1-4C-alkylpiperazino or morpholino group,

R4 is hydrogen, 1-4C-alkyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy, 1-4C-alkoxycarbonyl, trifluoromethyl, amino, mono- or di-1-4C-alkylamino, 1-4C-alkylcarbonylamino, 1-4C-alkoxycarbonylamino or 1-4C-alkoxy-1-4C-alkoxycarbonylamino,

R5 is hydrogen, 1-4C-alkyl or 1-4C-alkoxy and

X is O (oxygen) or NH,

and the salts of these compounds.

6. Compounds of the formula 1-1 according to claim 4, in which

R1 is 1-4C-alkyl,

R2 is 1-4C-alkyl,

R3 is carboxyl, 1-4C-alkoxycarbonyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy-1-4C-alkyl or the group -CO-NR31R32,

where

R31 is hydrogen, 1-4C-alkyl, hydroxy-1-4C-alkyl or 1-4C-alkoxy-1-4C-alkyl and

R32 is hydrogen or 1-4C-alkyl,

R4 is 1-4C-alkyl or 1-4C-alkylcarbonylamino,

R5 is 1-4C-alkyl and

X is O (oxygen) or NH,

and their salts.

7. Compounds of the formula 1-1 according to claim 4, in which

R1 is 1-4C-alkyl,

R2 is 1-4C-alkyl,

R3 is carboxyl, 1-4C-alkoxycarbonyl, hydroxy-1-4C-alkyl, 1-4C-alkoxy-1-4C-alkyl or the group -CO-NR31R32,

where

R31 is hydrogen, 1-4C-alkyl or hydroxy-1-4C-alkyl and

R32 is hydrogen or 1-4C-alkyl,

R4 is 1-4C-alkyl,

R5 is 1-4C-alkyl and

X is O (oxygen) or NH,

and their salts.

8. A compound according to claim 4 or 5 or 6 or 7, in which X is NH, or a salt of such compound.
9. A medicament comprising a compound as claimed in claim 1 and/or a pharmacologically acceptable salt thereof together with customary pharmaceutical auxiliaries and/or excipients.
10. The use of compounds as claimed in claim 1 and their pharmacologically acceptable salts for the prevention and treatment of gastrointestinal disorders.
11. The use of compounds as claimed in claim 1 and their pharmacologically acceptable salts for the production of medicaments which are suited for the prevention and treatment of gastrointestinal disorders.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP2004/050135

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D487/04 A61K31/495 A61P1/04

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, BEILSTEIN Data, PAJ

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EP 0 204 285 A (FUJISAWA PHARMACEUTICAL CO) 10 December 1986 (1986-12-10) claim 1	1-10
Y	KAMINSKI J J ET AL: "ANTIULCER AGENTS 2. GASTRIC ANTISECRETORY, CYTOPROTECTIVE, AND METABOLIC PROPERTIES OF SUBSTITUTED IMIDAZOU1,2-APYRIDINES AND ANALOGUES" JOURNAL OF MEDICINAL CHEMISTRY, AMERICAN CHEMICAL SOCIETY. WASHINGTON, US, vol. 30, no. 11, 1987, pages 2031-2046, XP002008621 ISSN: 0022-2623 examples 7,67,84; table III ----- -/--	1-10

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the international search

2 July 2004

Date of mailing of the international search report

19/07/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Baston, E

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP2004/050135

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 99/28322 A (DAHLSTROEM MIKAEL ;AMIN KOSRAT (SE); ASTRA AB (SE); NORDBERG PETER) 10 June 1999 (1999-06-10) cited in the application claims 1,10	1-10
Y	----- WERMUTH ET AL: "The Practise of Medicinal Chemistry" 1996, PRACTICE OF MEDICINAL CHEMISTRY, XX, XX, PAGE(S) 203-237 , XP002190259 the whole document	1-10
A	----- PATENT ABSTRACTS OF JAPAN vol. 1996, no. 01, 31 January 1996 (1996-01-31) & JP 07 242666 A (FUJISAWA PHARMACEUT CO LTD), 19 September 1995 (1995-09-19) abstract	1-10
A	----- WO 02/060492 A (CYTOPIA PTY LTD ;BURNS CHRISTOPHER JOHN (AU); WILKS ANDREW FREDERI) 8 August 2002 (2002-08-08) the whole document	1-10

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP2004/050135

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 0204285	A	10-12-1986	AT 71625 T	15-02-1992
			AU 593802 B2	22-02-1990
			AU 5834586 A	11-12-1986
			CA 1257264 A1	11-07-1989
			CN 86104313 A	04-03-1987
			DE 3683403 D1	27-02-1992
			DK 250386 A	05-12-1986
			EP 0204285 A1	10-12-1986
			ES 8800943 A1	16-02-1988
			FI 862210 A	05-12-1986
			GR 861379 A1	28-08-1986
			HU 40798 A2	27-02-1987
			JP 62016483 A	24-01-1987
			NO 862208 A	05-12-1986
			US 4725601 A	16-02-1988
			ZA 8603805 A	29-04-1987
			EP 0228006 A1	08-07-1987
			JP 62187471 A	15-08-1987
			US 4782055 A	01-11-1988
WO 9928322	A	10-06-1999	AT 233263 T	15-03-2003
			AU 752187 B2	12-09-2002
			AU 1356599 A	16-06-1999
			BR 9814755 A	03-10-2000
			CA 2311798 A1	10-06-1999
			CN 1142933 C	24-03-2004
			CZ 20001947 A3	15-11-2000
			DE 69811735 D1	03-04-2003
			DE 69811735 T2	18-12-2003
			DK 1042324 T3	05-05-2003
			EE 200000315 A	15-10-2001
			EP 1042324 A1	11-10-2000
			ES 2191356 T3	01-09-2003
			HK 1030216 A1	20-06-2003
			HU 0100601 A2	28-09-2001
			ID 24730 A	03-08-2000
			JP 2001525322 T	11-12-2001
			NO 20002721 A	28-07-2000
			NZ 504355 A	21-12-2001
			PL 340920 A1	12-03-2001
			PT 1042324 T	30-06-2003
			WO 9928322 A1	10-06-1999
			SI 1042324 T1	30-06-2003
			SK 6742000 A3	12-03-2001
			TR 200001530 T2	21-11-2000
			TW 515798 B	01-01-2003
			US 6518270 B1	11-02-2003
			ZA 9810468 A	21-05-1999
JP 07242666	A	19-09-1995	NONE	
WO 02060492	A	08-08-2002	WO 02060492 A1	08-08-2002
			CA 2436487 A1	08-08-2002
			EP 1363702 A1	26-11-2003
			US 2004102455 A1	27-05-2004